

FEniCS Course

Lecture 0: Introduction to FEM

Contributors
Anders Logg




FENICS
PROJECT

What is FEM?

The finite element method is a framework and a recipe for discretization of differential equations

- Ordinary differential equations
- Partial differential equations
- Integral equations
- A recipe for discretization of PDE
- PDE $\rightarrow Ax = b$
- Different bases, stabilization, error control, adaptivity

The FEM cookbook



The PDE (i)

Consider Poisson's equation, the Hello World of partial differential equations:

$$\begin{aligned} -\Delta u &= f && \text{in } \Omega \\ u &= u_0 && \text{on } \partial\Omega \end{aligned}$$

Poisson's equation arises in numerous applications:

- heat conduction, electrostatics, diffusion of substances, twisting of elastic rods, inviscid fluid flow, water waves, magnetostatics, . . .
- as part of numerical splitting strategies for more complicated systems of PDEs, in particular the Navier–Stokes equations

From PDE (i) to variational problem (ii)

The simple recipe is: multiply the PDE by a test function v and integrate over Ω :

$$-\int_{\Omega} (\Delta u)v \, dx = \int_{\Omega} f v \, dx$$

Then integrate by parts and set $v = 0$ on the Dirichlet boundary:

$$-\int_{\Omega} (\Delta u)v \, dx = \int_{\Omega} \nabla u \cdot \nabla v \, dx - \underbrace{\int_{\partial\Omega} \frac{\partial u}{\partial n} v \, ds}_{=0}$$

We find that:

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} f v \, dx$$

The variational problem (ii)

Find $u \in V$ such that

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} f v \, dx$$

for all $v \in \hat{V}$

The trial space V and the test space \hat{V} are (here) given by

$$V = \{v \in H^1(\Omega) : v = u_0 \text{ on } \partial\Omega\}$$

$$\hat{V} = \{v \in H^1(\Omega) : v = 0 \text{ on } \partial\Omega\}$$

From continuous (ii) to discrete (iii) problem

We approximate the continuous variational problem with a discrete variational problem posed on finite dimensional subspaces of V and \hat{V} :

$$V_h \subset V$$

$$\hat{V}_h \subset \hat{V}$$

Find $u_h \in V_h \subset V$ such that

$$\int_{\Omega} \nabla u_h \cdot \nabla v \, dx = \int_{\Omega} f v \, dx$$

for all $v \in \hat{V}_h \subset \hat{V}$

From discrete variational problem (iii) to discrete system of equations (iv)

Choose a basis for the discrete function space:

$$V_h = \text{span} \{ \phi_j \}_{j=1}^N$$

Make an ansatz for the discrete solution:

$$u_h = \sum_{j=1}^N U_j \phi_j$$

Test against the basis functions:

$$\int_{\Omega} \underbrace{\nabla \left(\sum_{j=1}^N U_j \phi_j \right)}_{u_h} \cdot \nabla \phi_i \, dx = \int_{\Omega} f \phi_i \, dx$$

From discrete variational problem (iii) to discrete system of equations (iv), contd.

Rearrange to get:

$$\sum_{j=1}^N U_j \underbrace{\int_{\Omega} \nabla \phi_j \cdot \nabla \phi_i \, dx}_{A_{ij}} = \underbrace{\int_{\Omega} f \phi_i \, dx}_{b_i}$$

A linear system of equations:

$$AU = b$$

where

$$A_{ij} = \int_{\Omega} \nabla \phi_j \cdot \nabla \phi_i \, dx \quad (1)$$

$$b_i = \int_{\Omega} f \phi_i \, dx \quad (2)$$

The canonical abstract problem

(i) Partial differential equation:

$$\mathcal{A}u = f \quad \text{in } \Omega$$

(ii) Continuous variational problem: find $u \in V$ such that

$$a(u, v) = L(v) \quad \text{for all } v \in \hat{V}$$

(iii) Discrete variational problem: find $u_h \in V_h \subset V$ such that

$$a(u_h, v) = L(v) \quad \text{for all } v \in \hat{V}_h$$

(iv) Discrete system of equations for $u_h = \sum_{j=1}^N U_j \phi_j$:

$$AU = b$$

$$A_{ij} = a(\phi_j, \phi_i)$$


$$b_i = L(\phi_i)$$

Important topics

- *How to choose V_h ?*
- *How to compute A and b*
- *How to solve $AU = b$?*
- *How large is the error $e = u - u_h$?*
- Extensions to nonlinear problems

How to choose V_h

Finite element function spaces




The finite element definition (Ciarlet 1975)

A finite element is a triple $(T, \mathcal{V}, \mathcal{L})$, where


- the domain T is a bounded, closed subset of \mathbb{R}^d (for $d = 1, 2, 3, \dots$) with nonempty interior and piecewise smooth boundary
- the space $\mathcal{V} = \mathcal{V}(T)$ is a finite dimensional function space on T of dimension n
- the set of degrees of freedom (nodes) $\mathcal{L} = \{\ell_1, \ell_2, \dots, \ell_n\}$ is a basis for the dual space \mathcal{V}' ; that is, the space of bounded linear functionals on \mathcal{V}

The finite element definition (Ciarlet 1975)

T



\mathcal{V}



\mathcal{L}

$$v(\bar{x})$$


$$v(\bar{x}) \cdot n$$

$$\int_T v(x)w(x) dx$$



The linear Lagrange element: $(T, \mathcal{V}, \mathcal{L})$

- T is a line, triangle or tetrahedron
- \mathcal{V} is the first-degree polynomials on T
- \mathcal{L} is point evaluation at the vertices

The linear Lagrange element: \mathcal{L}




The linear Lagrange element: \mathcal{V}_h




The quadratic Lagrange element: $(T, \mathcal{V}, \mathcal{L})$

- T is a line, triangle or tetrahedron
- \mathcal{V} is the second-degree polynomials on T
- \mathcal{L} is point evaluation at the vertices and edge midpoints

The quadratic Lagrange element: \mathcal{L}




The quadratic Lagrange element: \mathcal{V}_h



Families of elements



Families of elements



Computing the sparse matrix A

Naive assembly algorithm

$A = 0$

for $i = 1, \dots, N$

for $j = 1, \dots, N$

$$A_{ij} = a(\phi_j, \phi_i)$$

end for

end for

The element matrix

The global matrix A is defined by

$$A_{ij} = a(\phi_j, \phi_i)$$

The *element matrix* A_T is defined by

$$A_{T,ij} = a_T(\phi_j^T, \phi_i^T)$$

The assembly algorithm

$A = 0$

for $T \in \mathcal{T}$


 Compute the element matrix A_T

 Compute the local-to-global mapping ι_T

 Add A_T to A according to ι_T

end for

Adding the element matrix A_T



Solving $AU = b$

Direct methods

- Gaussian elimination
 - Requires $\sim \frac{2}{3}N^3$ operations
- LU factorization: $A = LU$
 - Solve requires $\sim \frac{2}{3}N^3$ operations
 - Reuse L and U for repeated solves
- Cholesky factorization: $A = LL^\top$
 - Works if A is symmetric and positive definite
 - Solve requires $\sim \frac{1}{3}N^3$ operations
 - Reuse L for repeated solves

Iterative methods

Krylov subspace methods

- GMRES (Generalized Minimal RESidual method)
- CG (Conjugate Gradient method)
 - Works if A is symmetric and positive definite
- BiCGSTAB, MINRES, TFQMR, ...

Multigrid methods

- GMG (Geometric MultiGrid)
- AMG (Algebraic MultiGrid)

Preconditioners


- ILU, ICC, SOR, AMG, Jacobi, block-Jacobi, additive Schwarz, ...

Which method should I use?

Rules of thumb

- Direct methods for small systems
- Iterative methods for large systems
- Break-even at ca 100–1000 degrees of freedom
- Use a symmetric method for a symmetric system
 - Cholesky factorization (direct)
 - CG (iterative)
- Use a multigrid preconditioner for Poisson-like systems
- GMRES with ILU preconditioning is a good default choice

Current timings (2012-01-20)



Homework!

- Install FEniCS 1.0.0!
- Download the FEniCS book!
- Visit the course web page!



`http://fenicsproject.org/`

`http://fenicsproject.org/pub/course/`

PS: Be alert and ready for the FEniCS challenge(s)...