

CHALMERS


FINITE ELEMENT CENTER



PREPRINT 2004–16

Adaptive variational multiscale methods based on a posteriori error estimation: Duality techniques for elliptic problems

Mats G. Larson and Axel Målqvist



Chalmers Finite Element Center

CHALMERS UNIVERSITY OF TECHNOLOGY

Göteborg Sweden 2004

CHALMERS FINITE ELEMENT CENTER

Preprint 2004–16

Adaptive variational multiscale methods based on a posteriori error estimation: Duality techniques for elliptic problems

Mats G. Larson and Axel Målqvist



CHALMERS

Chalmers Finite Element Center
Chalmers University of Technology
SE-412 96 Göteborg Sweden
Göteborg, May 2004

Adaptive variational multiscale methods based on a posteriori error estimation: Duality techniques for elliptic problems

Mats G. Larson and Axel Målqvist

NO 2004–16

ISSN 1404–4382

Chalmers Finite Element Center
Chalmers University of Technology
SE–412 96 Göteborg
Sweden

Telephone: +46 (0)31 772 1000

Fax: +46 (0)31 772 3595

www.phi.chalmers.se

Printed in Sweden
Chalmers University of Technology
Göteborg, Sweden 2004

Adaptive variational multiscale methods based on a posteriori error estimation: Duality techniques for elliptic problems

Mats G. Larson ^{*} Axel Målqvist [†]

May 18, 2004

Abstract

The variational multiscale method (VMM) provides a general framework for construction of multiscale finite element methods. In this paper we propose a method for parallel solution of the fine scale problem based on localized Dirichlet problems which are solved numerically. Next we present a posteriori error representation formulas for VMM which relates the error in linear functionals to the discretization errors, resolution and size of patches in the localized problems, in the fine scale approximation. These formulas are derived by using duality techniques. Based on the a posteriori error representation formula we propose an adaptive VMM with automatic tuning of the critical parameters. We primary study elliptic second order partial differential equations with highly oscillating coefficients or localized singularities.

1 Introduction

Many problems in science and engineering involve models of physical systems on many scales. For instance, models of materials with microstructure such as composites and flow in porous media. In such problems it is in general not feasible to seek for a numerical solution which resolves all scales. Instead we may seek to develop algorithms based on a suitable combination with a global problem capturing the main features of the solution and localized problems which resolves the fine scales. Since the fine scale problems are localized the computation on the fine scales is parallel in nature.

^{*}Corresponding author, Department of Computational Mathematics, Chalmers University of Technology, Göteborg, S-412 96, Sweden, mgl@math.chalmers.se

[†]Department Computational Mathematics, Chalmers University of Technology, Göteborg, S-412 96, Sweden, axel@math.chalmers.se

Previous work. The Variational Multiscale Method (VMM) is a general framework for derivation of basic multiscale method in a variational context, see Hughes [7] and [9]. The basic idea is to decompose the solution into fine and coarse scale contributions, solve the fine scale equation in terms of the residual of the coarse scale solution, and finally eliminate the fine scale solution from the coarse scale equation. This procedure leads to a modified coarse scale equation where the modification accounts for the effect of fine scale behavior on the coarse scales. In practice it is necessary to approximate the fine scale equation to make the method realistic. In several works various ways of analytical modeling are investigated often based on bubbles or element Green's functions, see Oberai and Pinsky, [11] and Arbogast [1]. In [6] Hou and Wu present a different approach. Here the fine scale equations are solved numerically on a finer mesh. The fine scale equations are solved inside coarse elements and are thus totally decoupled.

New contributions. In this work we present a simple technique for numerical approximation of the fine scale equation in the variational multiscale method. The basic idea is to split the fine scale residual into localized contributions using a partition of unity and solving corresponding decoupled localized problems on patches with homogeneous Dirichlet boundary conditions. The fine scale solution is approximated by the sum $U_f = \sum_i U_{f,i}$ of the solutions $U_{f,i}$ to the localized problems. The accuracy of U_f depends on the fine scale mesh size h and the size of the patches. We note that the fine scale computation is naturally parallel.

To optimize performance we seek to construct an adaptive algorithm for automatic control of the coarse mesh size H , the fine mesh size h , and the size of patches. Our algorithm is based on the following a posteriori estimate of the error $e = u - U_c - U_f$ for the Poisson equation with variable coefficients a :

$$(e, \psi) = \sum_{i \in \mathcal{C}} (\varphi_i R(U_c), \phi_f) + \sum_{i \in \mathcal{F}} ((\varphi_i R(U_c), \phi_f)_{\omega_i} - a(U_{f,i}, \phi_f)_{\omega_i}), \quad (1.1)$$

where $\psi \in H^{-1}(\Omega)$, \mathcal{C} refers to nodes where no local problems have been solved, \mathcal{F} to nodes where local problems are solved, U_c is the coarse scale solution, $U = U_c + U_f$, $R(U) = f + \nabla \cdot a \nabla U$ is the residual, and ϕ_f is the fine scale part of a dual solution driven by ψ .

If no fine scale equations are solved only we obtain the first term in the estimate. The second term relates the fine scale mesh parameter h to the patch size ω_i on which the local problems are solved. We have derived a similar estimate for the error in energy norm, see [10].

The framework is fairly general and may be extended to other types of multiscale methods, for instance, based on localized Neumann problems.

Outline. First we introduce the model problem and the variational multiscale formulation of this problem, we also discuss the split of the coarse and fine scale spaces. In the following section we present a posteriori estimates of the error. These results leads

to an adaptive algorithm. We present numerical results and finally we present concluding remarks and suggestions on future work.

2 The Variational Multiscale Method

2.1 Model Problem

We study the Poisson equation with a highly oscillating coefficient a and homogeneous Dirichlet boundary conditions: find $u \in H_0^1(\Omega)$ such that

$$-\nabla \cdot a \nabla u = f \quad \text{in } \Omega, \quad (2.1)$$

where Ω is a polygonal domain in \mathbf{R}^d , $d = 1, 2$, or 3 with boundary Γ , $f \in H^{-1}(\Omega)$, and $a \in L^\infty(\Omega)$ such that $a(x) \geq \alpha_0 > 0$ for all $x \in \Omega$. The variational form of (2.1) reads: find $u \in \mathcal{V} = H_0^1(\Omega)$ such that

$$a(u, v) = (f, v) \quad \text{for all } v \in \mathcal{V}, \quad (2.2)$$

with the bilinear form

$$a(u, v) = (a \nabla u, \nabla v) \quad (2.3)$$

for all $u, v \in \mathcal{V}$.

2.2 The Variational Multiscale Method

We employ the variational multiscale scale formulation, proposed by Hughes see [7, 9] for an overview, and introduce a coarse and a fine scale in the problem. We choose two spaces $\mathcal{V}_c \subset \mathcal{V}$ and $\mathcal{V}_f \subset \mathcal{V}$ such that

$$\mathcal{V} = \mathcal{V}_c \oplus \mathcal{V}_f. \quad (2.4)$$

Then we may pose (2.2) in the following way: find $u_c \in \mathcal{V}_c$ and $u_f \in \mathcal{V}_f$ such that

$$\begin{aligned} a(u_c, v_c) + a(u_f, v_c) &= (f, v_c) \quad \text{for all } v_c \in \mathcal{V}_c, \\ a(u_c, v_f) + a(u_f, v_f) &= (f, v_f) \quad \text{for all } v_f \in \mathcal{V}_f. \end{aligned} \quad (2.5)$$

Introducing the residual $R : \mathcal{V} \rightarrow \mathcal{V}'$ defined by

$$(R(v), w) = (f, w) - a(v, w) \quad \text{for all } w \in \mathcal{V}, \quad (2.6)$$

the fine scale equation takes the form: find $u_f \in \mathcal{V}_f$ such that

$$a(u_f, v_f) = (R(u_c), v_f) \quad \text{for all } v_f \in \mathcal{V}_f. \quad (2.7)$$

Thus the fine scale solution is driven by the residual of the coarse scale solution. Denoting the solution u_f to (2.7) by $u_f = TR(u_c)$ we get the modified coarse scale problem

$$a(u_c, v_c) + a(TR(u_c), v_c) = (f, v_c) \quad \text{for all } v_c \in \mathcal{V}_c. \quad (2.8)$$

Here the second term on the left hand side accounts for the effects of fine scales on the coarse scales.

2.3 A VMM Based on Localized Dirichlet Problems

We introduce a partition $\mathcal{K} = \{K\}$ of the domain Ω into shape regular elements K of diameter H_K and we let \mathcal{N} be the set of nodes. Further we let \mathcal{V}_c be the space of continuous piecewise polynomials of degree p defined on \mathcal{K} .

We shall now construct an algorithm which approximates the fine scale equation by a set of decoupled localized problems. We begin by writing $u_f = \sum_{i \in \mathcal{N}} u_{f,i}$ where

$$a(u_{f,i}, v_f) = (\varphi_i R(u_c), v_f) \quad \text{for all } v_f \in \mathcal{V}_f, \quad (2.9)$$

and $\{\varphi_i\}_{i \in \mathcal{N}}$ is the set of Lagrange basis functions in \mathcal{V}_c . Note that $\{\varphi_i\}_{i \in \mathcal{N}}$ is a partition of unity with support on the elements sharing the node i . We call the set of elements with one corner in node i a mesh star in node i and denote it S_1^i . Thus functions $u_{f,i}$ correspond to the fine scale effects created by the localized residuals $\varphi_i R(u_c)$. Introducing this expansion of u_f in the right hand side of the fine scale equation (2.5) and get: find $u_c \in \mathcal{V}_c$ and $u_f = \sum_{i \in \mathcal{N}} u_{f,i} \in \mathcal{V}_f$ such that

$$\begin{aligned} a(u_c, v_c) + a(u_f, v_c) &= (f, v_c) \quad \text{for all } v_c \in \mathcal{V}_c, \\ a(u_{f,i}, v_f) &= (\varphi_i R(u_c), v_f) \quad \text{for all } v_f \in \mathcal{V}_f \text{ and } i \in \mathcal{N}. \end{aligned} \quad (2.10)$$

We use this fact to construct a finite element method for solving (2.10) approximately in two steps.

- For each coarse node we define a patch ω_i such that $\text{supp}(\varphi_i) \subset \omega_i \subset \Omega$. We denote the boundary of ω_i by $\partial\omega_i$.
- On these patches we define piecewise polynomial spaces $\mathcal{V}_f^h(\omega_i)$ with respect to a fine mesh with mesh function $h = h(x)$ defined as a piecewise constant function on the fine mesh. Functions in $\mathcal{V}_f^h(\omega_i)$ are equal to zero on $\partial\omega_i$.

The resulting method reads: find $U_c \in \mathcal{V}_c$ and $U_f = \sum_i^n U_{f,i}$ where $U_{f,i} \in \mathcal{V}_f^h(\omega_i)$ such that

$$\begin{aligned} a(U_c, v_c) + a(U_f, v_c) &= (f, v_c) \quad \text{for all } v_c \in \mathcal{V}_c, \\ a(U_{f,i}, v_f) &= (\varphi_i R(U_c), v_f) \quad \text{for all } v_f \in \mathcal{V}_f^h(\omega_i) \text{ and } i \in \mathcal{N}. \end{aligned} \quad (2.11)$$

Since the functions in the local finite element spaces $\mathcal{V}_f^h(\omega_i)$ are equal to zero on $\partial\omega_i$, U_f and therefore $U = U_c + U_f$ will be continuous.

Remark 2.1 For problems with multiscale phenomena on a part of the domain it is not necessary to solve local problems for all coarse nodes. We let $\mathcal{C} \subset \mathcal{N}$ refer to nodes where no local problems are solved and $\mathcal{F} \subset \mathcal{N}$ refer to nodes where local problems are solved. Obviously $\mathcal{C} \cup \mathcal{F} = \mathcal{N}$. We let $U_{f,i} = 0$ for $i \in \mathcal{C}$.

Remark 2.2 The choice of the subdomains ω_i is crucial for the method. We introduce a notation for extended stars of many layers of coarse elements recursively in the following way. The extended mesh star $S_L^i = \cup_{j \in S_{L-1}^i} S_1^j$ for $L > 1$. We refer to L as layers, see Figure 1.

In Figure (2) we plot solutions to localized fine scale problems $U_{f,i}$ on different patches.




Figure 1: Two (left) and one (right) layer stars.




Figure 2: The fine scale solution $U_{f,i}$ for different patches.




Figure 3: HB-function and WHB-function with two Jacobi iterations.

2.4 Subspaces

The choice of the fine scale space \mathcal{V}_f can be done in different ways. In a paper by Aksoylu and Holst [4] three suggestions are made.

Hierarchical basis method. The first and perhaps easiest approach is to let $\mathcal{V}_f = \{v \in \mathcal{V} : v(x_j) = 0, j = \mathcal{N}\}$, where $\{x_i\}_{i \in \mathcal{N}}$ are the coarse mesh nodes. When \mathcal{V}_f is discretized by the standard piecewise polynomials on the fine mesh this means that the fine scale base functions will have support on fine scale stars.

BPX preconditioner. The second approach is to let \mathcal{V}_f be $L^2(\Omega)$ orthogonal to \mathcal{V}_c . In this case we will have global support for the fine scale base functions but for the discretized space we have rapid decay outside fine mesh stars.

Wavelet modified hierarchical basis method. The third choice is a mix of the other two. The fine scale space \mathcal{V}_f is defined as an approximate $L^2(\Omega)$ orthogonal version of the Hierarchical basis method. We let $Q_c^a v \in \mathcal{V}_c$ be an approximate solution (a small number of Jacobi iterations) to

$$(Q_c^a v, w) = (v, w), \quad \text{for all } w \in \mathcal{V}_c. \quad (2.12)$$

and define the Wavelet modified hierarchical basis function associated with the hierarchical basis function φ_{HB} to be,

$$\varphi_{WHB} = (I - Q_c^a)\varphi_{HB}, \quad (2.13)$$

see Figure 3.

For an extended description of these methods see [3, 4, 2]. In this paper we focus on the WHB method.

3 A Posteriori Error Estimates

3.1 The Dual Problem

To derive a posteriori error estimates of the error in a given linear functional (e, ψ) with $e = u - U$ and $\psi \in H^{-1}(\Omega)$ a given weight. We introduce the following dual problem: find

$\phi \in \mathcal{V}$ such that

$$a(v, \phi) = (v, \psi) \quad \text{for all } v \in \mathcal{V}. \quad (3.1)$$

In the VMM setting this yields: find $\phi_c \in \mathcal{V}_c$ and $\phi_f \in \mathcal{V}_f$ such that

$$\begin{aligned} a(v_c, \phi_c) + a(v_c, \phi_f) &= (v_c, \psi), \quad \text{for all } v_c \in \mathcal{V}_c, \\ a(v_f, \phi_f) + a(v_f, \phi_c) &= (v_f, \psi), \quad \text{for all } v_f \in \mathcal{V}_f. \end{aligned} \quad (3.2)$$

3.2 Error Representation Formula

We now derive an error representation formula involving both the coarse scale error $e_c = u_c - U_c$ and the fine scale error $e_f = \sum_{i \in \mathcal{N}} e_{f,i} := \sum_{i \in \mathcal{N}} (u_{f,i} - U_{f,i})$ that arises from using our finite element method (2.11).

We use the dual problem (3.2) to derive an a posteriori error estimate for a linear functional of the error $e = e_c + e_f$. If we subtract the coarse part of equation (2.11) from the coarse part of equation (2.10) we get the Galerkin orthogonality,

$$a(e_c, v_c) + a(e_f, v_c) = 0 \quad \text{for all } v_c \in \mathcal{V}_c. \quad (3.3)$$

The same argument on the fine scale equation gives for $i \in \mathcal{F}$,

$$a(e_{f,i}, v_f) = (f, \varphi_i v_f) - a(e_c, \varphi_i v_f), \quad \text{for all } v_f \in \mathcal{V}_f^h(\omega_i). \quad (3.4)$$

We are now ready to state an error representation formula.

Theorem 3.1 *If $\psi \in H^{-1}(\Omega)$ then,*

$$(e, \psi) = \sum_{i \in \mathcal{C}} (\varphi_i R(U_c), \phi_f) + \sum_{i \in \mathcal{F}} ((\varphi_i R(U_c), \phi_f)_{\omega_i} - a(U_{f,i}, \phi_f)_{\omega_i}). \quad (3.5)$$

Proof. Starting from the definition of the dual problem and letting $v = e = u - U_c - U_f$ we get

$$(e, \psi) = a(e, \phi) \quad (3.6)$$

$$= a(e, \phi_f) \quad (3.7)$$

$$= a(u - U_c, \phi_f) - a(U_f, \phi_f) \quad (3.8)$$

$$= (R(U_c), \phi_f) - a(U_f, \phi_f) \quad (3.9)$$

$$= (R(U_c), \phi_f) - \sum_{i \in \mathcal{F}} a(U_{f,i}, \phi_f) \quad (3.10)$$

$$= \sum_{i \in \mathcal{C}} (\varphi_i R(U_c), \phi_f) + \sum_{i \in \mathcal{F}} ((\varphi_i R(U_c), \phi_f)_{\omega_i} - a(U_{f,i}, \phi_f)_{\omega_i}). \quad (3.11)$$

which proves the theorem. \square

Since equation (2.11) holds we can subtract functions $v_{f,i}^h \in \mathcal{V}_f^h(\omega_i)$ where $i \in \mathcal{F}$ from equation (3.11). For example we choose $v_{f,i}^h = \pi_{h,i}\phi_f$, where $\pi_{h,i}\phi_f$ is the Scott-Zhang interpolant of ϕ_f onto $\mathcal{V}_f^h(\omega_i)$ to get

$$(e, \psi) = \sum_{i \in \mathcal{C}} (\varphi_i R(U_c), \phi_f) \quad (3.12)$$

$$+ \sum_{i \in \mathcal{F}} ((\varphi_i R(U_c), \phi_f - \pi_{h,i}\phi_f)_{\omega_i} - a(U_{f,i}, \phi_f - \pi_{h,i}\phi_f)_{\omega_i}).$$

Remark 3.1 Since the dual problem defined in equation (3.2) is equally hard to solve as the primal problem we need to solve it numerically as well. Normally it would not be sufficient to solve the dual problem with the same accuracy as the primal due to the Galerkin Orthogonality. However in this setting things are a bit different. Calculating ϕ_f with the same accuracy as U_f or even with lower accuracy will not result in an error equal to zero. The important thing is to only store the part of ϕ_f with support on ω_i when calculating term i in the sum of equation (3.5). The entire function ϕ_f might be hard to store in the memory of the computer.

4 Adaptive Algorithm

We use the error representation formula in Theorem 3.1 to construct an adaptive algorithm. We remember the result,

$$(e, \psi) = \sum_{i \in \mathcal{C}} (\varphi_i R(U_c), \phi_f) + \sum_{i \in \mathcal{F}} ((\varphi_i R(U_c), \phi_f)_{\omega_i} - a(U_{f,i}, \phi_f)_{\omega_i}). \quad (4.1)$$

The first sum of the error representation formula is very similar to what we would get from using standard Galerkin on the coarse mesh. The function $\phi_f = \phi - \phi_c \sim H\nabla\phi$ which is exactly what we would expect. For the second sum we have an extra orthogonality namely that from equation (3.12). We have $\phi_f - \pi_{h,i}\phi_f \sim h\nabla\phi$ if the patches $\omega_i = \Omega$ i.e. we get the fine scale convergence. But in practice the patches are much smaller so we end up somewhere in between h and H convergence. To sum up this discussion there are three parameters of interest that need to be considered in an adaptive algorithm, H , h , and the size of the patches.

Adaptive Algorithm.

- Start with no nodes in \mathcal{F} .
- Calculate the primal U_c .
- Calculate the dual solution locally ϕ_f with low accuracy for all coarse nodes. (ψ_f does not need to be solved very accurately to point out the correct nodes for local calculations.)




Figure 4: Geometry (left) and Reference solution (right).

- Calculate the contributions to the error for each coarse node, $C_i = (\varphi_i R(U_c), \phi_f)$.
- Solve local problems where C_i is large to get a new U_c .
- Calculate C_i and $F_i = ((\varphi_i R(U_c), \phi_f)_{\omega_i} - a(U_{f,i}, \phi_f)_{\omega_i})$, for large values in C_i solve more local problems and for large values in F_i either increase the number of layers or decrease the fine scale mesh size h for local problem i . Stop if the desired tolerance is reached or calculate a new U_c and new error indicators etc. .

5 Numerical Examples

We solve two dimensional model problems with linear base functions defined on a uniform triangular mesh.

Example 1. In this example we demonstrate how we can get highly improves accuracy in one part of the domain by choosing the load in the dual problem ψ equal to the indicator function for this domain. We consider the unit square with a crack in the form of a plus sign on which the solution is forced to be zero, see Figure 5 (left). We let ψ be equal to one in the lower left quadrant (marked with a thin lattice in the figure) and zero in the rest of the domain. To the right in Figure 5 we see a reference solution to the Poisson equation with $a = f = 1$ and homogeneous Dirichlet boundary conditions on this geometry. The idea is to use the adaptive algorithm to choose which areas that needs to be solved with higher accuracy. In Figure 5 we plot the dual solution, with ψ chosen as described above, to the left and the fine scale part of the dual solution to the right. After two iterations in the adaptive algorithm we see clearly that local problems have only nodes in the lower left corner. In Figure 5 the small circles refers to fine scale problems solved with two layer stars and the bigger circles refers to fine scale problems solved three layer stars. The improvement in the solution after two iteration in the adaptive algorithm is very clear. In Figure 5 we compare the standard Galerkin solution and the adaptive solution to a




Figure 5: Dual solution (left) and fine scale part of the dual solution (right) calculated with $\psi = I_{\{0 \leq x, y \leq 0.5\}}$.




Figure 6: Local problems solved with two and three layer stars.




Figure 7: Galerkin error (left) and adaptive variational multiscale method error (right).




Figure 8: The coefficient $a = 1$ on the white parts and $a = 0.05$ on the lattice (left) and reference solution on this geometry (right).

reference solution. We see how the error in the lower left quadrant is much smaller but the error in the rest of the domain is very similar to the standard Galerkin error.

Example 2. Next we turn our attention to a model problem with oscillating coefficient a in a part of the domain, see Figure 5. In this example we choose $f = \psi = 1$ which makes the primal and the dual equivalent. In Figure 5 we note that the adaptive algorithm automatically picks nodes in the left part of the domain for local problems to increase accuracy. In the first example we want to refine a certain part of the domain and therefore we choose ψ in order to do so, here we want good accuracy on the whole domain and the adaptive algorithm chooses where to refine automatically. Again we compare standard Galerkin and our solution to a reference solution calculated on a finer mesh. The result can be seen in Figure 5. Again we see a nice improvement compared to the standard Galerkin error. The choice $\psi = 1$ indicates control of the mean of the error over the domain.




Figure 9: The dots marks coarse nodes where local problems have been solved.




Figure 10: Standard Galerkin error (left) and the error using adaptivity (right).

6 Conclusions and Future Work

We have presented a method for parallel solution of the fine scale equations in the variational multiscale method based on solution of localized Dirichlet problems on patches and developed an a posteriori error analysis for the method. Based on the estimates we design a basic adaptive algorithm for automatic tuning of the critical parameters: resolution and size of patches in the fine scale problems. It is also possible to decide whether a fine scale computation is necessary or not and thus the proposed scheme may be combined with a standard adaptive algorithm on the coarse scales. The method is thus very general in nature and may be applied to any problem where adaptivity is needed.

In this paper we have focused on two scales in two spatial dimensions. A natural extension would be to solve three dimensional problems with multiple scales. It is also natural to extend this theory to other equations modeling for instance flow and materials. We also intend to study non-linear and time dependent equations using this approach.

References

- [1] T. Arbogast and S. L. Bryant, A two-scale numerical subgrid technique for waterflood simulations, SPE J., Dec. 2002, pp 446-457.
- [2] B. Aksoylu, S. Bond, and M. Holst *An odyssey into local refinement and multilevel preconditioning III: Implementation of numerical experiments*, SIAM J. Sci. Comput., 25(2003), 478-498.
- [3] B. Aksoylu and M. Holst *An odyssey into local refinement and multilevel preconditioning I: Optimality of the BPX preconditioner*, SIAM J. Numer. Anal. (2002) in review
- [4] B. Aksoylu and M. Holst *An odyssey into local refinement and multilevel preconditioning II: stabilizing hierarchical basis methods*, SIAM J. Numer. Anal. in review
- [5] S. C. Brenner and L. R. Scott, *The mathematical theory of finite element methods*, Springer Verlag, 1994.
- [6] T. Y. Hou and X.-H. Wu, *A multiscale finite element method for elliptic problems in composite materials and porous media*, J. Comput. Phys. 134 (1997) 169-189.
- [7] T. J.R. Hughes, *Multiscale phenomena: Green's functions, the Dirichlet-to-Neumann formulation, subgrid scale models, bubbles and the origins of stabilized methods*, Comput. Methods Appl. Mech. Engrg. 127 (1995) 387-401.
- [8] T. J.R. Hughes, *The continuous Galerkin method is locally conservative*, J. Comput. Phys., Vol. 163(2) (2000) 467-488.

- [9] T. J.R. Hughes, G. R. Feijóo, L. Mazzei and Jean-Baptiste Quincy, *The variational multiscale method - a paradigm for computational mechanics*, Comput. Methods Appl. Mech. Engrg. 166 (1998) 3-24.
- [10] M. G. Larson and A. Målqvist, *Adaptive Variational Multiscale Methods Based on A Posteriori Error Estimates: Energy Estimates for Elliptic Problems*, in preparation
- [11] A. A. Oberai and P. M. Pinsky, *A multiscale finite element method for the Helmholtz equation*, Comput. Methods Appl. Mech. Engrg. 154 (1998) 281-297.

Chalmers Finite Element Center Preprints

- 2003–01** *A hybrid method for elastic waves*
Larisa Beilina
- 2003–02** *Application of the local nonobtuse tetrahedral refinement techniques near Fichera-like corners*
L. Beilina, S. Korotov and M. Křížek
- 2003–03** *Nitsche’s method for coupling non-matching meshes in fluid-structure vibration problems*
Peter Hansbo and Joakim Hermansson
- 2003–04** *Crouzeix–Raviart and Raviart–Thomas elements for acoustic fluid–structure interaction*
Joakim Hermansson
- 2003–05** *Smoothing properties and approximation of time derivatives in multistep backward difference methods for linear parabolic equations*
Yubin Yan
- 2003–06** *Postprocessing the finite element method for semilinear parabolic problems*
Yubin Yan
- 2003–07** *The finite element method for a linear stochastic parabolic partial differential equation driven by additive noise*
Yubin Yan
- 2003–08** *A finite element method for a nonlinear stochastic parabolic equation*
Yubin Yan
- 2003–09** *A finite element method for the simulation of strong and weak discontinuities in elasticity*
Anita Hansbo and Peter Hansbo
- 2003–10** *Generalized Green’s functions and the effective domain of influence*
Donald Estep, Michael Holst, and Mats G. Larson
- 2003–11** *Adaptive finite element/difference method for inverse elastic scattering waves*
Larisa Beilina
- 2003–12** *A Lagrange multiplier method for the finite element solution of elliptic domain decomposition problems using non-matching meshes*
Peter Hansbo, Carlo Lovadina, Ilaria Perugia, and Giancarlo Sangalli
- 2003–13** *A reduced P^1 –discontinuous Galerkin method*
R. Becker, E. Burman, P. Hansbo, and M.G. Larson
- 2003–14** *Nitsche’s method combined with space–time finite elements for ALE fluid–structure interaction problems*
Peter Hansbo, Joakim Hermansson, and Thomas Svedberg
- 2003–15** *Stabilized Crouzeix–Raviart element for the Darcy–Stokes problem*
Erik Burman and Peter Hansbo

- 2003–16** *Edge stabilization for the generalized Stokes problem: a continuous interior penalty method*
Erik Burman and Peter Hansbo
- 2003–17** *A conservative flux for the continuous Galerkin method based on discontinuous enrichment*
Mats G. Larson and A. Jonas Niklasson
- 2003–18** *CAD-to-CAE integration through automated model simplification and adaptive modelling*
K.Y. Lee, M.A. Price, C.G. Armstrong, M.G. Larson, and K. Samuelsson
- 2003–19** *Multi-adaptive time integration*
Anders Logg
- 2003–20** *Adaptive computational methods for parabolic problems*
Kenneth Eriksson, Claes Johnson, and Anders Logg
- 2003–21** *The FEniCS project*
T. Dupont, J. Hoffman, C. Johnson, R.C. Kirby, M.G. Larson, A. Logg, and R. Scott
- 2003–22** *Adaptive finite element methods for LES: Computation of the mean drag coefficient in a turbulent flow around a surface mounted cube using adaptive mesh refinement*
Johan Hoffman
- 2003–23** *Adaptive DNS/LES: a new agenda in CFD*
Johan Hoffman and Claes Johnson
- 2003–24** *Multiscale convergence and reiterated homogenization of parabolic problem*
Anders Holmbom, Nils Svanstedt, and Niklas Wellander
- 2003–25** *On the relationship between some weak compactnesses with different numbers of scales*
Anders Holmbom, Jeanette Silfver, Nils Svanstedt, and Niklas Wellander
- 2003–26** *A posteriori error estimation in computational inverse scattering*
Larisa Beilina and Claes Johnson
- 2004–01** *Computability and adaptivity in CFD*
Johan Hoffman och Claes Johnson
- 2004–02** *Interpolation estimates for piecewise smooth functions in one dimension*
Anders Logg
- 2004–03** *Estimates of derivatives and jumps across element boundaries for multi-adaptive Galerkin solutions of ODEs*
Anders Logg
- 2004–04** *Multi-adaptive Galerkin methods for ODEs III: Existence and stability*
Anders Logg
- 2004–05** *Multi-adaptive Galerkin methods for ODEs IV: A priori error estimates*
Anders Logg
- 2004–06** *A stabilized non-conforming finite element method for incompressible flow*
Erik Burman and Peter Hansbo

- 2004–07** *On the uniqueness of weak solutions of Navier-Stokes equations: Remarks on a Clay Institute prize problem*
Johan Hoffman and Claes Johnson
- 2004–08** *A new approach to computational turbulence modeling*
Johan Hoffman and Claes Johnson
- 2004–09** *A posteriori error analysis of the boundary penalty method*
Kenneth Eriksson, Mats G. Larson, and Axel Målqvist
- 2004–10** *A posteriori error analysis of stabilized finite element approximations of the helmholtz equation on unstructured grids*
Mats G. Larson and Axel Målqvist
- 2004–11** *Adaptive variational multiscale methods based on a posteriori error estimation*
Mats G. Larson and Axel Målqvist
- 2004–12** *Multi-adaptive Galerkin methods for ODEs V: Stiff problems*
Johan Jansson and Anders Logg
- 2004–13** *Algorithms for multi-adaptive time-stepping*
Johan Jansson and Anders Logg
- 2004–14** *Simulation of mechanical systems with individual time steps*
Johan Jansson and Anders Logg
- 2004–15** *Computational modeling of dynamical systems*
Johan Jansson, Claes Johnson, and Anders Logg
- 2004–16** *Adaptive variational multiscale methods based on a posteriori error estimation: Duality techniques for elliptic problems*
Mats G. Larson and Axel Målqvist

These preprints can be obtained from

www.phi.chalmers.se/preprints